Bedrock Geologic Map of the Salem (Iowa) 7.5' Quadrangle



1:24,000

0 0.5 1 2 3 4 Miles

0 0.5 1 2 3 4 Kilometers

MN
GN
0°50'
15 MILS
0°58'
17 MILS

UTM GRID AND 2019 MAGNETIC NORTH DECLINATION AT CENTER OF SHEET

Adjacent 7.5' Quadrangles

LOCKRIDGE EAST, IOWA	MOUNT PLEASANT, IOWA	NEW LONDON, IOWA
HILLSBORO, IOWA	SALEM, IOWA	LOWELL, IOWA
FARMINGTON, IOWA	DONNELLSON, IOWA	WEST POINT, IOWA

Location Map



Legend

CENOZOIC

QUATERNARY SYSTEM

Qu – **Qu - Undifferentiated Unconsolidated Sediments** - Consists of loamy soils developed in loess, glacial till, and colluvium of variable thickness, and alluvial clay, silt, sand, and gravel. The total thickness of the Quaternary deposits varies between 0 and 18 m (0 – 60 ft), but can be as much as 50 m (165 ft) thick in the eastern part of the mapping area. This unit is shown only on the cross-section, not on the map.

PALEOZOIC

CARBONIFEROUS SYSTEM

PENNSYLVANIAN SUBSYSTEM

Pcl – **Pcl - Shale and Sandstone** (lower Cherokee Group) Lower-Middle Pennsylvanian. Pennsylvanian units occur as erosional outliers reaching a thickness of up to 13 m (45 ft) within the mapping area. This unit consists of shale/mudstone, light to medium gray, part silty to sandy and fine to medium quartz sandstone, rarely conglomeratic. Some shales are carbonaceous to phosphatic with minor coal. No outcrops of this unit were identified in the mapping area.

MISSISSIPPIAN SUBSYSTEM

Mpsl – **Mpsl – Limestone, Sandstone, and Dolomite** (Pella or "St. Louis" formations) Middle-Upper Mississippian, Meramecian-lower Chesterian. This map unit ranges between 9 and 18 m (30 – 60 ft) thick and reaches a maximum thickness of 29 m (95 ft) in the mapping area. It is dominated by limestone, sandstone, dolomitic limestone, and dolomite with minor shale and chert. Limestones of the Pella Formation are typically sub-lithographic with scattered to abundant fossils, primarily brachiopods, echinoderms, and ostracods. The "St. Louis" Formation is dominated by limestone, sandy limestone, sandstone, and dolomite, variably cherty. The limestone facies of this unit can be fossiliferous with brachiopods, echinoderms, and several varieties of coral while the dolomitic facies typically exhibit fossil molds. Some fossils are silicified. Sandstones of the "St. Louis" Formation are typically very fine to medium quartz sandstones that are poorly to moderately cemented with calcite or quartz. The lower portion of the "St. Louis" Formation is commonly gray to dark brown dolomite, locally brecciated and sandy, with rare fossils. This mapping unit dominates the bedrock surface in the mapping area and is overlain by Quaternary sediments or Pennsylvanian outliers. No outcrops were identified in the mapping area although excellent exposures of the lower portion of this mapping unit were observed in a small active quarry in the extreme southeast corner of the mapping area.

Mws – **Mws – Shale, Dolomite, and Limestone** (Warsaw Formation) Upper Osagean. The Warsaw Formation varies in thickness reaching a maximum thickness of approximately 17 m (55 ft). This unit can generally be divided into two major lithologic groupings, a lower argillaceous dolomite sequence and an upper shale dominated sequence. The upper shale is typically light to medium gray, silty, and variably dolomitic with minor chert, sand, and sparse quartz geodes. The lower dolomite, sometimes referred to as the "geode beds," is argillaceous to shaly, with scattered to abundant quartz geodes. Minor limestone units occur locally as thin, lensatic beds with crinoidal packstone/grainstone fabrics. Brachiopods, echinoderm debris, and bryozoans are found throughout this mapping unit, although these fossils are more common in the carbonate lithologies. This unit exhibits wide variability, leaving only the upper shale or lower dolomite in place, suggesting strong erosional unconformities above and below this mapping unit. Outcrops of this unit were not observed in the mapping area.

Mkeo – **Mkeo – Limestone, Dolomite, Chert, and Shale** (Keokuk Formation) Upper Osagean. The Keokuk Formation typically ranges from 12 to 23 m (40 – 75 ft) in thickness in the mapping area. This unit is dominated by tan to gray interbedded skeletal limestones displaying packstone/grainstone fabrics. Nodular to bedded chert, in part fossiliferous, is common in the lower half of the sequence. Dolomite, variably argillaceous, and thin shales also occur throughout the unit. The unit displays multiple hardground surfaces and bone beds with scattered to abundant fish debris, the most prominent of these serves as a marker bed at the base of the formation (sometimes referred to as the Burlington-Keokuk or B-K bone bed). Brachiopods, crinoids, bryozoans, solitary corals, and fish bones and teeth occur throughout this unit as both abraded debris and partly articulated specimens. Molds of sponge spicules are noted in the dolomite facies. Traces of glauconite and locally abundant geodes were also observed within the mapping area. Outcrops of this unit were not observed in the mapping area.

Mb – **Mb – Limestone, Dolomite, and Chert** (Burlington Formation) Lower Osagean. The Burlington Formation can be up to 29 m (95 ft) thick in the mapping area. This unit is subdivided into three members (in ascending order: the Dolbee Creek, Haight Creek, and Cedar Fork), characterized by distinct lithologic groupings. The Dolbee Creek Member is dominated by white to tan skeletal limestone displaying packstone/grainstone fabrics and nodular to bedded chert. The Haight Creek Member is characterized by dolomite with an intermittent unit of skeletal limestone (sometimes referred to as the "middle grainstone") and thick beds of chert. A glauconite-rich zone marks the lower contact between the Dolbee Creek and can be used as a regional marker bed. Fossil molds are also present in the dolomite facies. The Cedar Fork Member is a pure white crinoidal packstone limestone unit which is usually differentiated from the packstones of the overlying Keokuk Formation by its white appearance. Occasional fish debris and glauconite are also observed in this member. Outcrops of the Burlington Formation were not found in the mapping area.

Mk – **Mk – Dolomite, Limestone, and Siltstone** (Kinderhookian formations) Lower Mississippian. The Kinderhookian sequence ranges in thickness from 5 to 15 m (20 – 50 ft) with a maximum thickness of 27 m (90 ft) in the mapping area. This unit comprises three formations (in ascending order: the McCraney, Prospect Hill, and Wassonville), characterized by distinct lithologic groupings. The McCraney Formation is composed of alternating beds of sparsely fossiliferous, sub-lithographic limestone and dark brown, unfossiliferous dolomite generating a unique "zebra striped" appearance in outcrop. A basal oolite is locally present. The Prospect Hill Formation is a light to medium gray, dolomitic siltstone that grades to shale in some locations. This unit is often laminated with vertical and horizontal burrow fabrics and faint cross stratified bedforms. Fossils are rare to absent although fossil molds are locally abundant. The Wassonville Formation, now including the former Starr's Cave Formation as the basal member, consists of massive dolomite that is variably cherty grading into dolomitic limestone lower in the section. The basal Starr's Cave Member is a fossiliferous limestone with packstone/grainstone fabrics and is commonly oolitic. Crinoids (partly articulated) are the dominant fossil type of the Starr's Cave Member. A diverse assemblage of brachiopods are present with lesser amounts of blastoids, starfish, corals, bryozoans, and trilobites reported.

DEVONIAN SYSTEM

Der – **Der – Siltstone and Shale** (English River Formation) Upper Devonian, Famennian. The English River Formation is up to 6 m (20 ft) within the mapping area. This unit is dominated by gray to olive-green siltstone with apparent bioturbated fabrics. Bivalves and brachiopods are common, especially in the upper beds, with scattered to abundant fossil molds as well. This unit only appears in the cross-section, not on the map.

Dss – **Dss – Shale** (Saverton Shale Formation) Upper Devonian, Famennian. The Saverton Shale Formation can be up to 30 m (100 ft) within the mapping area. This unit is dominated by green-gray shale, commonly burrowed with sparse to absent macrofossils. This unit only appears in the cross-section, not on the map.

OTHER FEATURES

- **New drill holes for this map project**
- **New geophysics data point**
- **Bedrock outcrops**
- **IGS GEOSAM data points** – records available at www.iowageologicalsurvey.org
- **Quarries**
- **Incorporated towns or cities**
- **Roads**
- W7036 **Wells used for geologic cross-section**
- **Bedrock Hillshade**- shades of gray show the bedrock surface as it would be illuminated by an artificial light source from the NW direction

Stratigraphic Column

System[1]	Subsystem[1]	Series[1]	Stage[2]	Lithostratigraphic Unit	Map Symbol	Lithology	Thickness (in feet)
Carboniferous	Pennsylvanian	Lower	Morrowan?	lower Cherokee Group	Pcl		<45
Carboniferous	Mississippian	Middle: Upper	Chesterian	Pella or "St. Louis" fms.	Mpsl		<95
Carboniferous	Mississippian	Middle	Meramecian	Pella or "St. Louis" fms.	Mpsl		<95
Carboniferous	Mississippian	Middle	Osagean	Warsaw Fm.	Mws		<55
Carboniferous	Mississippian	Middle	Osagean	Keokuk Fm.	Mkeo		40-75
Carboniferous	Mississippian	Lower	Osagean	Burlington Fm.	Mb		<95
Carboniferous	Mississippian	Lower	Kinderhookian	Kinderhookian fms.	Mk		20-50
Devonian		Upper	Famennian	English River Fm.	Der		<20
Devonian		Upper	Famennian	Saverton Shale Fm.	Dss		<100

[1]Global stratigraphic units
[2]Regional stratigraphic units

Lithologies: dolomite; dolomitic shale; fossiliferous packstone; limestone; lithographic limestone; sandstone; sandy limestone; shale; siltstone

Symbols: geodes; chert; oolitic; argillaceous zone; breccia; unconformity

Geologic Cross-Section A-A'



Bedrock Geologic Map of the Salem 7.5' Quadrangle, Henry and Lee Counties, Iowa

Iowa Geological Survey
Open File Map OFM-19-3
June 2019

Ryan Clark, Stephanie Tassier-Surine, and Phil Kerr

Iowa Geological Survey, IIHR-Hydroscience & Engineering, University of Iowa, Iowa City, Iowa

Iowa Geological Survey, Keith Schilling, State Geologist

Supported in part by the U.S. Geological Survey
Cooperative Agreement Number G18AC00194
National Cooperative Geologic Mapping Program (STATEMAP)
This work was partially funded by a National Science Foundation Award:
Improving Undergraduate STEM Education: GP-IMPACT-1600429

ACKNOWLEDGMENTS

Special thanks to Don Smith of Cessford Construction, Co. (Oldcastle Materials) for allowing us to access quarries in and around the mapping area during field activities. University of Iowa (UI) Department of Earth and Environmental Sciences student Travis Maher and Cornell College student Gabi Hiatt prepared well cutting samples for stratigraphic logging. New subsurface geologic data was generated by UI students Megan Koch and Alethea Kapolas, by producing descriptive logs of water well drilling samples. Megan Koch and UI student Allison Kasick also helped with well locations and data management. Jason Vogelgesang of the Iowa Geological Survey (IGS) helped with geophysical data acquisition. Thanks also to Rick Langel of the IGS for managing the Iowa geologic sampling database (GeoSam). Brian Witzke (IGS – retired), Illinois State University Professor of Geology James "Jed" Day, UI Assistant Professors of Geology Brad Cramer and Emily Finzel are all thanked for their help with the stratigraphic research components stemming from this mapping project. Casey Kohrt and Chris Kahle of the Iowa Department of Natural Resources provided GIS technical help. Bill Bunker, Ray Anderson, Bob McKay, and Brian Witzke (IGS – retired) provided valued background information concerning the bedrock topography, geology, and Mississippian and Devonian stratigraphy of the area. Administrative support was provided by Suzanne Doershuk, Melissa Eckrich, Teresa Gaffey, Carmen Langel, and Rosemary Tiwari.

Introduction to the Bedrock Geology of the Salem 7.5' Quadrangle, Henry and Lee Counties, Iowa

The Salem Quadrangle lies within the Southern Iowa Drift Plain landform region (Prior, 1991). The map area is dominated by loess-mantled till plains in the uplands, and fine to coarse grained alluvial deposits within the Skunk River and its tributaries. This area hosts glacial deposits of Pre-Illinoian age (ranging from 0.5 to 2.6 million years ago). The thickness of Quaternary materials overlying the bedrock surface varies widely across the quadrangle ranging from 0 to 18 m (0 – 60 ft), reaching a maximum thickness of 50 m (165 ft) within the bedrock valley in the northeastern part of the mapping area. Shallow bedrock information from the soil surveys of Henry and Lee counties (Seaholm, 1985; and Lockridge, 1979) was used for identifying potential bedrock outcrop locations during field mapping activities. Bedrock outcrops were not observed during field activities, however according to the soil survey maps, outcrops exist along tributaries of the Skunk River, and in areas of the southeastern part of the quadrangle. There is one operational quarry in the extreme southeastern corner of the quadrangle that was observed during field mapping activities, as well as one abandoned quarry that was not accessible. Subsurface information was mostly derived from the analysis of water well cutting samples reposited at the Iowa Geological Survey (IGS). Lithologic and stratigraphic information from these samples are stored in the online GeoSam database of the IGS. Geologic information from one quarry and more than 250 private and public wells within the Salem Quadrangle and the surrounding area were used for bedrock geologic mapping purposes.

The Mississippian System (now Subsystem) was historically proposed for the succession of strata exposed in the Mississippi River Valley between Burlington, Iowa and southern Illinois. Therefore, the bedrock exposures in southeastern Iowa take on a special significance as they comprise part of the historic "body stratotype" on which the concept of the Mississippian System was defined and based (Witzke et al., 2002). The Mississippian had been primarily a North American chronostratigraphic label roughly synonymous with the Lower Carboniferous of the Old World. After approval by the Subcommission on Carboniferous Stratigraphy in 1999 and ratification by the International Union of Geological Sciences and the International Commission on Stratigraphy in 2000, the Carboniferous System was officially subdivided into the lower and upper subsystems, the Mississippian and Pennsylvanian, respectively. As such, the Mississippian now has meaning and application as a major subdivision of geologic time not only in North America, but as a globally defined subsystem. The bedrock strata seen in the Salem Quadrangle and surrounding area provide a significant historic reference for the Mississippian as a whole.

The conundrum that is the Mississippian in Iowa has been the subject of curiosity for many previous workers. Owen (1852) and Hall (1857) were the first to recognize that the abundant bedrock exposures in southeastern Iowa likely correlated with those observed farther down the Mississippi River Valley. Decades later, Van Tuyl (1923) took on the ambitious task of correlating all of the Mississippian units across Iowa. Many of their lithologic interpretations were valuable; however, the correlations were, and continue to be, subject to revision as later workers attempted to piece the Mississippian into the global stratigraphic framework. Harris and Parker (1964) provided inspirational insights into the structural context of southeastern Iowa by identifying a series of northwest-southeast trending anticlines that were later found to be superimposed on the larger northeast-southwest trending structural feature known as the Mississippi Arch (Witzke et al., 1990). Many questions remain regarding the stratigraphic correlations within the Mississippian such as whether the "St. Louis" Formation in Iowa truly belongs in the St. Louis Formation or whether some of the upper members be reassigned to the Ste. Genevieve Formation; whether the Prospect Hill Formation is an offshoot of the Hannibal Formation of Missouri and Illinois; and whether the McCraney Formation is correlative to the McCraney in Illinois or if it should become a new stratigraphic interval (as proposed by Witzke et al., 2002). In an effort to address the question regarding the "St. Louis" Formation, detrital zircon analyses from sandstone samples collected near the mapping area were processed with the help of Emily Finzel (Assistant Professor of Geology at the University of Iowa (UI)). The geochronologic data provided by the detrital zircon analyses were not able to differentiate the sandstone units within the "St. Louis" Formation, however, further study of the geochemistry and lithology of these sandstones may provide the evidence needed to identify whether these units belong in the St. Louis proper or in the Ste. Genevieve. Clarification of the issue regarding the Prospect Hill and McCraney formations is being carried out with the help of Brad Cramer (Assistant Professor of Geology at the UI), Brittany Stolfus (UI student), and James "Jed" Day (Professor of Geology at Illinois State University). Samples collected from locations southeast of the mapping area as well as at other locations in eastern Illinois and northeastern Missouri for conodonts and carbon isotopes have provided valuable bio- and chemostratigraphic information. Preliminary results suggest that the Prospect Hill and McCraney formations in Iowa may correlate with the Hannibal Formation of Missouri and Illinois. Further study will commence with additional sampling of surface exposures as well as core samples. Rectifying the questions posed by Witzke et al., 2002, may now become attainable.

Although the Mississippian bedrock in southeastern Iowa is no longer a widely used aquifer due to low yields and locally poor water quality, many of the bedrock units are highly desirable sources of aggregate, thus necessitating the continued effort to gain a better understanding of the local and regional stratigraphic characteristics and relationships of the Mississippian Subsystem in southeastern Iowa.

The mapping area consists of bedrock of the Mississippian Subsystem from late Kinderhookian to early Chesterian (about 355 – 330 million years ago) and Devonian strata of Famennian age (about 370 – 360 million years ago) (Ogg et al., 2008). Famennian strata are represented by brown, organic rich shales of the Grassy Creek Formation overlain by gray-green silty shales of the Saverton Shale Formation and capped by the English River Formation siltstone. The thick shale packages represent major transgressive-regressive cycles of deposition in a stratified seaway (Witzke, 1987). Kinderhookian strata represent a sequence of interbedded carbonates and siltstones that unconformably underlie the Burlington Formation (early Osagean) within the mapping area. The Burlington, Keokuk, and Warsaw formations (collectively the Augusta Group of Witzke et al., 2010) represent a relatively conformable package of marine rocks deposited during the Osagean transgressive-regressive (T-R) cycle. Interpreted as part of the central middle shelf of the Osagean sea that transgressed toward the northwest and the Transcontinental Arch, the Burlington Formation rocks were deposited across a vast subtidal epicontinental shelf that stretched from Illinois and Iowa into central Kansas and Oklahoma (Lane, 1978; Witzke et al., 1990). The Keokuk and Warsaw formations represent the regressive phase of the Osagean T-R cycle punctuated by a stark unconformity below the overlying Pella or "St. Louis" formations, regionally displaying up to 40 m (130 ft) of erosional relief (Witzke et al., 2002). For a more detailed description of the lithologic units and further discussion of mapping methodologies, please refer to the accompanying Summary Report.

References

Hall, J., 1857: Observations upon the Carboniferous limestones of the Mississippi Valley. American Journal of Science, v. 23, p. 187-203.
Harris Jr., S.E. and Parker, M.C., 1964: Stratigraphy of the Osage Series in Southeastern Iowa, Iowa Geological Survey Report of Investigation No. 1. 52 p., 17 figs, 7 plates.
Lane, H.R., 1978: The Burlington Shelf (Mississippian, north-central United States). Geologica et Palaeontologica v. 12, p. 165-176.
Lockridge, L.D., 1979: Soil Survey of Lee County, Iowa. U.S. Department of Agriculture, Soil Conservation Service, 188 p., 85 map sheets.
Ogg, J.G., Ogg, G., and Gradstein, F.M., 2008: The Concise Geologic Time Scale, Cambridge University Press, 177 p.
Owen, D.D., 1852: Report of a Geological Survey of Wisconsin, Iowa, and Minnesota; and incidentally of a portion of Nebraska territory. Lippincott, Gambo and Co., 638 p.
Prior, J.C., 1991: Landforms of Iowa. Univ. of Iowa Press, Iowa City, 154 p.
Seaholm, J.E., 1985: Soil Survey of Henry County, Iowa. U.S. Department of Agriculture, Soil Conservation Service, 259 p., 63 map sheets.
Van Tuyl, F.M., 1923: The stratigraphy of the Mississippian formations of Iowa. Iowa Geological Survey, Annual Report, v. 30, p. 33-359.
Witzke, B.J., 1987: Models for circulation patterns in epicontinental seas applied to Paleozoic facies of North American craton. Paleoceanography, v. 2, p. 229-248.
Witzke, B.J., Anderson, R.R., and Pope, J.P., 2010: Bedrock Geologic Map of Iowa, 1:500,000. Iowa Geological and Water Survey Open File Map OFM-10-1.
Witzke, B.J., Bunker, B.J., Anderson, R.R., Artz, J.A., and Tassier-Surine, S.A., 2002: Pleistocene, Mississippian, & Devonian Stratigraphy of the Burlington, Iowa, Area. Iowa Geological Survey Guidebook No. 23, 137 p.
Witzke, B.J., McKay, R.M., Bunker, B.J., and Woodson, F.J., 1990: Stratigraphy and Paleoenvironments of Mississippian Strata in Keokuk and Washington Counties, Southeast Iowa. Iowa Geological Survey Guidebook No. 10, 105 p.

Base map from USGS Salem 7.5' Digital Raster Graphic (IGS GIS file IA_Salem_USGS_topo.tif) which was scanned and modified from the Salem 7.5' Topographic Quadrangle map, published by The US Geological Survey in 2015
Land elevation contours (10' interval). Bedrock topography raster created internally for this map project.

Iowa Geological Survey digital cartographic file Salem_BedrockGeology.mxd, version 6/30/19 (ArcGIS 10.5)
Map projection and coordinate system based on Universal Transverse Mercator (UTM) Zone 15N, datum NAD83.

The map and cross-section are based on interpretations of the best available information at the time of mapping. Map interpretations are not a substitute for detailed site specific studies.

Research supported by the U. S. Geological Survey, National Cooperative Geologic Mapping Program, under USGS award numberG18AC00194. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U. S. Government.